

Record: 1

Title: The Diagnosis and Treatment of Glaucoma.

Authors: Schuster AK; Department of Ophthalmology, University Medical Center Mainz; Private Institute of Applied Ophthalmology Berlin; Department of Ophthalmology, University Hospital Cologne.
Erb C
Hoffmann EM
Dietlein T
Pfeiffer N

Source: Deutsches Arzteblatt international [Dtsch Arztebl Int] 2020 Mar 27; Vol. 117 (13), pp. 225-234.

Publication Type: Journal Article; Review

Language: English

Journal Info: *Publisher:* Deutscher Ärzte-Verlag *Country of Publication:* Germany
NLM ID: 101475967 *Publication Model:* Print *Cited Medium:* Internet
ISSN: 1866-0452 (Electronic) *Linking ISSN:* 18660452 *NLM ISO Abbreviation:* Dtsch Arztebl Int *Subsets:* MEDLINE

Imprint Name(s): *Original Publication:* Cologne : Deutscher Ärzte-Verlag

MeSH Terms: Glaucoma/*diagnosis
Glaucoma/*therapy
Europe/epidemiology ; Glaucoma/epidemiology ; Humans

Abstract: Background: Glaucoma is a group of chronically progressive disorders of the optic nerve. In this article, we present the epidemiology of and risk factors for glaucoma, as well as the diagnostic work-up and treatment options.

Methods: This review is based on pertinent publications retrieved by a selective search in Medline and the Cochrane Library, supplemented by further articles chosen by the authors.

Results: In Europe, the prevalence of glaucoma is 2.93% among persons aged 40 to 80 years. The prevalence rises with age, reaching 10% in persons over 90 years old. The available diagnostic methods include ophthalmoscopy, tonometry, perimetry, and imaging techniques. The treatment of glaucoma is focused on lowering the intraocular pressure with topical drugs, laser therapy, and glaucoma surgery. In patients with manifest glaucoma, lowering the intraocular pressure prevents the progression of visual field defects, with a number needed to treat of 7.

Conclusion: The diagnostic evaluation of glaucoma rests on multiple pillars, all of which must be considered for establishing the diagnosis and defining the desired target pressure: these are, among others, the intraocular pressure and ocular function and morphology. Individually tailored pressure-lowering treatment should be evaluated in regularly scheduled follow-up visits for assessment of function and morphology and adjusted as necessary to minimize the risk of progression.

- References:** Isr J Med Sci. 1972 Aug-Sep;8(8):1362-5. (PMID: 4265110)
Br J Ophthalmol. 2013 Feb;97(2):196-200. (PMID: 23203707)
Invest Ophthalmol Vis Sci. 2005 Apr;46(4):1275-9. (PMID: 15790890)
Ophthalmology. 2010 Sep;117(9):1705-12. (PMID: 20591487)
J Ophthalmol. 2017;2017:2190347. (PMID: 29085672)
Ophthalmology. 2016 Mar;123(3):532-41. (PMID: 26746594)
JAMA. 2014 May 14;311(18):1901-11. (PMID: 24825645)
Ophthalmology. 2017 Oct;124(10):1466-1474. (PMID: 28549518)
Cochrane Database Syst Rev. 2015 Nov 06;(11):CD006259. (PMID: 26545176)
Cochrane Database Syst Rev. 2014 Feb 15;(2):CD007059. (PMID: 24532137)
Cochrane Database Syst Rev. 2007 Oct 17;(4):CD003167. (PMID: 17943780)
BMJ. 2005 Jul 16;331(7509):134. (PMID: 15994659)
PLoS One. 2016 Oct 20;11(10):e0162229. (PMID: 27764086)
Lancet. 2017 Nov 11;390(10108):2183-2193. (PMID: 28577860)
Ophthalmol Eye Dis. 2015 Oct 08;7:21-33. (PMID: 26483611)
Clin Exp Ophthalmol. 2015 Jan-Feb;43(1):40-6. (PMID: 24811050)
Acta Ophthalmol. 2019 Feb;97(1):60-67. (PMID: 29575745)
Ophthalmology. 2013 Jun;120(6):1120-6. (PMID: 23415421)
Ophthalmology. 1994 Oct;101(10):1662-7. (PMID: 7936564)
Surv Ophthalmol. 1999 Jan-Feb;43(4):293-320. (PMID: 10025513)
Prog Retin Eye Res. 2000 Jan;19(1):1-40. (PMID: 10614679)
PLoS One. 2015 Feb 23;10(2):e0115625. (PMID: 25705901)
Br J Ophthalmol. 1996 Dec;80(12):1068-72. (PMID: 9059272)
Surv Ophthalmol. 2008 Nov;53 Suppl1:S93-105. (PMID: 19038628)
Ophthalmologe. 2015 May;112(5):402-9. (PMID: 25930182)
PLoS One. 2015 Oct 01;10(10):e0138825. (PMID: 26425846)
PLoS One. 2018 Jan 12;13(1):e0191185. (PMID: 29329338)
Prog Brain Res. 2015;221:233-41. (PMID: 26518081)
Dtsch Arztebl Int. 2017 Mar 24;114(12):204-210. (PMID: 28407842)
Ophthalmology. 2018 May;125(5):650-663. (PMID: 29477688)
Acta Ophthalmol. 2012 Jun;90(4):316-21. (PMID: 20626719)
PLoS One. 2014 Aug 19;9(8):e102972. (PMID: 25137059)
Invest Ophthalmol Vis Sci. 2011 Aug 29;52(9):6875-81. (PMID: 21715354)
Ophthalmology. 2016 Apr;123(4):771-82. (PMID: 26795295)
Acta Ophthalmol. 2013 Dec;91(8):e612-9. (PMID: 23710686)
Invest Ophthalmol Vis Sci. 2003 Sep;44(9):3783-9. (PMID: 12939292)
J Glaucoma. 2009 Mar;18(3):253-61. (PMID: 19295383)
Invest Ophthalmol Vis Sci. 2017 Mar 1;58(3):1462-1468. (PMID: 28264102)
Lancet. 2016 Oct 1;388(10052):1389-1397. (PMID: 27707497)
Ophthalmology. 2014 Nov;121(11):2081-90. (PMID: 24974815)
Ophthalmol Ther. 2018 Jun;7(1):49-73. (PMID: 29725860)

Ophthalmology. 2004 Apr;111(4):651-64. (PMID: 15051195)
Acta Ophthalmol. 2015 Dec;93(8):e621-6. (PMID: 25913492)
Lancet. 2015 Apr 4;385(9975):1295-304. (PMID: 25533656)
PLoS One. 2015 Jul 06;10(7):e0131770. (PMID: 26147908)
Invest Ophthalmol Vis Sci. 2016 May 1;57(6):2452-62. (PMID: 27149695)
Acta Biomater. 2017 Jun;55:340-348. (PMID: 28323178)
PLoS One. 2017 Jul 25;12(7):e0181675. (PMID: 28742840)
PLoS One. 2017 Aug 29;12(8):e0183142. (PMID: 28850575)
Br J Ophthalmol. 2016 Jan;100(1):86-93. (PMID: 26286821)
Ophthalmology. 2016 Jan;123(1):129-40. (PMID: 26526633)
Arch Ophthalmol. 2002 Jun;120(6):701-13; discussion 829-30. (PMID: 12049574)
Acta Ophthalmol. 2019 Feb;97(1):e50-e56. (PMID: 30225863)
Clin Ophthalmol. 2017 Aug 29;11:1583-1600. (PMID: 28919702)
Br J Ophthalmol. 2018 May;102(5):575-585. (PMID: 29545417)
Expert Opin Drug Saf. 2008 Nov;7(6):795-9. (PMID: 18983225)
Graefes Arch Clin Exp Ophthalmol. 2018 Sep;256(9):1695-1702. (PMID: 29774418)
Am J Ophthalmol. 2014 Sep;158(3):615-27.e9. (PMID: 24879946)
J Glaucoma. 2014 Apr-May;23(4):246-53. (PMID: 23221903)
Ophthalmology. 2013 Aug;120(8):1541-5. (PMID: 23631945)
Ophthalmologica. 1957 Oct;134(4):221-42. (PMID: 13484216)
Arch Ophthalmol. 2002 Oct;120(10):1268-79. (PMID: 12365904)
Cochrane Database Syst Rev. 2018 Apr 25;4:CD009313. (PMID: 29694684)
Int Ophthalmol. 2018 Apr;38(2):429-441. (PMID: 28243830)
Klin Monbl Augenheilkd. 2017 Nov;234(11):1407-1422. (PMID: 28931182)
Prog Retin Eye Res. 2015 May;46:67-83. (PMID: 25619727)
Curr Opin Ophthalmol. 2017 Mar;28(2):154-160. (PMID: 27898469)

Entry Date(s): *Date Created:* 20200429 *Date Completed:* 20200908 *Latest Revision:* 20200908

Update Code: 20210210

PubMed Central ID: PMC7196841

DOI: 10.3238/arztebl.2020.0225

PMID: 32343668

Database: MEDLINE with Full Text

Record: 1

Title: Glaucoma Diagnosis and Treatment: The Role of the Ophthalmic Nurse.

Authors: Ney JJ

Source: Insight (American Society of Ophthalmic Registered Nurses) [Insight] 2016 Winter; Vol. 41 (1), pp. 13-17.

Publication Type: Journal Article; Review

Language: English

Journal Info: *Publisher:* Mosby For The American Society Of Ophthalmic Registered Nurses *Country of Publication:* United States *NLM ID:* 9111431
Publication Model: Print *Cited Medium:* Print *ISSN:* 1060-135X (Print)
Linking ISSN: 1060135X *NLM ISO Abbreviation:* Insight Subsets: Nursing

Imprint Name(s): *Publication:* St. Louis Mo : Mosby For The American Society Of Ophthalmic Registered Nurses
Original Publication: San Francisco, CA : The Society,

MeSH Terms: Nurse Specialists*
Nurse's Role*
Ophthalmology*
Glaucoma, Open-Angle/*diagnosis
Glaucoma, Open-Angle/*therapy
Nurse Practitioners/*statistics & numerical data
Diagnostic Techniques, Ophthalmological/nursing ; Humans ; Intraocular Pressure ; Workforce

Abstract: Glaucoma is one of the single largest causes of irreversible blindness. Glaucomatous vision loss is preventable with the appropriate diagnostic testing and treatment. Ophthalmic nurses play an important role in ensuring the success of glaucoma diagnosis and treatment.

Entry Date(s): *Date Created:* 20180920 *Date Completed:* 20181016 *Latest Revision:* 20181202

Update Code: 20210210

PMID: 30230735

Database: MEDLINE with Full Text

Record: 1

Title: Measurement of Retinal Changes in Primary Acute Angle Closure Glaucoma under Different Durations of Symptoms.

Authors: Zhu X; Department of Ophthalmology, Zhongnan Hospital of Wuhan University, Wuhan, China.
Zeng W; Department of Ophthalmology, Zhongnan Hospital of Wuhan University, Wuhan, China.
Wu S; Department of Ophthalmology, Zhongnan Hospital of Wuhan University, Wuhan, China.
Chen X; Department of Ophthalmology, Zhongnan Hospital of Wuhan University, Wuhan, China.
Zheng T; Department of Ophthalmology, Zhongnan Hospital of Wuhan University, Wuhan, China.
Ke M; Department of Ophthalmology, Zhongnan Hospital of Wuhan University, Wuhan, China.

Source: Journal of ophthalmology [J Ophthalmol] 2019 Dec 04; Vol. 2019, pp. 5409837. *Date of Electronic Publication:* 2019 Dec 04 (*Print Publication:* 2019).

Publication Type: Journal Article

Language: English

Journal Info: *Publisher:* Hindawi Publishing Corporation *Country of Publication:* United States *NLM ID:* 101524199 *Publication Model:* eCollection *Cited Medium:* Print *ISSN:* 2090-004X (Print) *Linking ISSN:* 2090004X *NLM ISO Abbreviation:* J Ophthalmol *Subsets:* PubMed not MEDLINE

Imprint Name(s): *Original Publication:* New York, NY : Hindawi Publishing Corporation

Abstract: **Purpose:** To investigate the changes of retinal nerve fiber layer (RNFL) in patients after an attack of primary acute angle closure glaucoma (PAACG) and to assess the impact of attack time on prognosis of retinal changes.

Design: cross-sectional study.

Methods: Twenty-six patients with unilateral PAACG attack and cataracts from 2017 to 2019 were enrolled. Eyes with PAACG attack time less than 1 day constituted the group A (n = 13), while eyes with PAACG attack time more than 1 day constituted the group B (n = 13). All patients received phacoemulsification and viscogoniosynechialysis after intraocular pressure (IOP) lowering. All patients underwent ophthalmic examinations including IOP, best-corrected visual acuity (BCVA), and visual field (VF). Optical coherence tomography angiography (OCTA) was used to obtain circumpapillary RNFL vessel density (cpVD). Spectral domain optical coherence tomography (SD-OCT) was used to examine the peripapillary RNFL and macular ganglion cell complex (GCC). All patients accepted 2 assessments before and 1 month after the procedure.

Results: The IOP of all patients recovered to normal (12.77 ± 2.65 mm

Hgvs. 12.77 ± 3.85 mmHg, $p = 0.834$) after the procedure. Patients in the group A had better BCVA improvement than those in the group B (1.32 ± 0.84 vs. 0.50 ± 0.21 , $p = 0.004$), as well as better mean defect (MD) values from VF (-3.65 ± 2.54 vs -16.05 ± 5.99 , $p < 0.001$). Compared with group B, patients in the group A had thicker macula (Fovea area: 255.00 ± 27.94 μ m vs. 203.92 ± 59.73 μ m, $p = 0.010$), thicker GCC (82.62 ± 8.76 μ m vs. 65.23 ± 18.56 μ m, $p = 0.005$), and thicker RNFL (105.08 ± 9.38 μ m vs. 77.69 ± 20.23 μ m, $p < 0.001$). Higher blood flow density in all-plexus peripapillary retina was observed in the group A eyes compared with group B (full sector: 0.56 ± 0.02 vs. 0.41 ± 0.07 , $p < 0.001$). In both groups, the association between average RNFL thickness and cpVD as well as MD values and pattern standard deviation (PSD) values from VF was stronger ($R^2 = 0.58, 0.60, -0.54$, respectively, all $p < 0.001$) than the association between GCC thickness and cpVD, as well as MD values and PSD values ($R^2 = 0.37, p = 0.001; R^2 = 0.37, p = 0.001; R^2 = -0.27, p = 0.007$).

Conclusion: Patients with attack time less than 1 day had better retinal thickness and all-plexus peripapillary retina blood flow density. Controlling the attack time could decrease retinal damage by PAACG. (Copyright © 2019 Xiaolu Zhu et al.)

- References:** Clin Ophthalmol. 2019 May 23;13:859-868. (PMID: 31190729)
 Acta Ophthalmol. 2016 May;94(3):e211-9. (PMID: 26132774)
 Graefes Arch Clin Exp Ophthalmol. 2015 Sep;253(9):1557-64. (PMID: 26255817)
 Int Ophthalmol. 2012 Dec;32(6):577-82. (PMID: 22847248)
 Graefes Arch Clin Exp Ophthalmol. 2017 May;255(5):1013-1018. (PMID: 28161831)
 Br J Ophthalmol. 1972 Mar;56(3):175-85. (PMID: 4624382)
 Aust N Z J Ophthalmol. 1988 Aug;16(3):245-50. (PMID: 3179051)
 PLoS One. 2017 Sep 15;12(9):e0184301. (PMID: 28915256)
 Br J Ophthalmol. 1977 Feb;61(2):126-32. (PMID: 843509)
 Prog Retin Eye Res. 1999 Jan;18(1):121-32. (PMID: 9920501)
 Invest Ophthalmol Vis Sci. 2016 Jul 1;57(9):OCT451-9. (PMID: 27409505)
 J Glaucoma. 2018 Apr;27(4):322-327. (PMID: 29419642)
 Br J Ophthalmol. 1994 Aug;78(8):643-8. (PMID: 7918293)
 Surv Ophthalmol. 1994 Jul-Aug;39(1):23-42. (PMID: 7974188)
 Med J Malaysia. 2013 Oct;68(5):410-4. (PMID: 24632871)
 PLoS One. 2012;7(6):e40352. (PMID: 22768284)
 Prog Brain Res. 2008;173:31-45. (PMID: 18929100)
 Jpn J Ophthalmol. 1995;39(2):180-6. (PMID: 8538076)
 Br J Ophthalmol. 2002 Feb;86(2):238-42. (PMID: 11815354)
 Am J Ophthalmol. 2017 May;177:106-115. (PMID: 28254626)
 Ophthalmology. 2014 Mar;121(3):750-8. (PMID: 24290800)
 Surv Ophthalmol. 1994 May;38 Suppl:S3-6. (PMID: 7940146)
 Arch Ophthalmol. 1977 Nov;95(11):1961-74. (PMID: 921573)

Br J Ophthalmol. 2006 Jan;90(1):14-6. (PMID: 16361658)

Jpn J Ophthalmol. 2014 May;58(3):252-60. (PMID: 24496567)

Invest Ophthalmol Vis Sci. 2019 Feb 1;60(2):552-558. (PMID: 30721302)

J Glaucoma. 2014 Feb;23(2):119-24. (PMID: 22895523)

Ophthalmology. 2016 Dec;123(12):2498-2508. (PMID: 27726964)

Entry Date(s): *Date Created:* 20191231 *Latest Revision:* 20201001

Update Code: 20210210

PubMed Central ID: PMC6914934

DOI: 10.1155/2019/5409837

PMID: 31885888

Database: MEDLINE with Full Text

Measurement of Retinal Changes in Primary Acute Angle Closure Glaucoma under Different Durations of Symptoms

1. Introduction

Purpose. To investigate the changes of retinal nerve fiber layer (RNFL) in patients after an attack of primary acute angle closure glaucoma (PAACG) and to assess the impact of attack time on prognosis of retinal changes. **Design.** cross-sectional study. **Methods.** Twenty-six patients with unilateral PAACG attack and cataracts from 2017 to 2019 were enrolled. Eyes with PAACG attack time less than 1 day constituted the group A (n = 13), while eyes with PAACG attack time more than 1 day constituted the group B (n = 13). All patients received phacoemulsification and viscosynovialysis after intraocular pressure (IOP) lowering. All patients underwent ophthalmic examinations including IOP, best-corrected visual acuity (BCVA), and visual field (VF). Optical coherence tomography angiography (OCTA) was used to obtain circumpapillary RNFL vessel density (cpVD). Spectral domain optical coherence tomography (SD-OCT) was used to examine the peripapillary RNFL and macular ganglion cell complex (GCC). All patients accepted 2 assessments before and 1 month after the procedure. **Results.** The IOP of all patients recovered to normal (12.77 ± 2.65 mm Hgvs. 12.77 ± 3.85 mmHg, $p=0.834$) after the procedure. Patients in the group A had better BCVA improvement than those in the group B (1.32 ± 0.84 vs. 0.50 ± 0.21 , $p=0.004$), as well as better mean defect (MD) values from VF (-3.65 ± 2.54 vs -16.05 ± 5.99 , $p<0.001$). Compared with group B, patients in the group A had thicker macula (Fovea area: 255.00 ± 27.94 μm vs. 203.92 ± 59.73 μm , $p=0.010$), thicker GCC (82.62 ± 8.76 μm vs. 65.23 ± 18.56 μm , $p=0.005$), and thicker RNFL (105.08 ± 9.38 μm vs. 77.69 ± 20.23 μm , $p<0.001$). Higher blood flow density in all-plexus peripapillary retina was observed in the group A eyes compared with group B (full sector: 0.56 ± 0.02 vs. 0.41 ± 0.07 , $p<0.001$). In both groups, the association between average RNFL thickness and cpVD as well as MD values and pattern standard deviation (PSD) values from VF was stronger ($R^{²} = 0.58, 0.60, -0.54$, respectively, all $p<0.001$) than the association between GCC thickness and cpVD, as well as MD values and PSD values ($R^{²} = 0.37$, $p=0.001$; $R^{²} = 0.37$, $p=0.001$; $R^{²} = -0.27$, $p=0.007$). **Conclusion.** Patients with attack time less than 1 day had better retinal thickness and all-plexus peripapillary retina blood flow density. Controlling the attack time could decrease retinal damage by PAACG.

Primary acute angle closure glaucoma (PAACG), an acute ocular hypertension disease that may lead to irreversible damage to the optic nerve and other ocular tissues, is one of the most common types of glaucoma, especially among Asians [1]. Ocular biometric characteristics of Asians result in higher prevalence in Asia [2]. Shallow anterior chamber depth and lens-related and retrolenticular mechanisms lead to overcrowding anterior

structure and decreased opening of trabecular meshwork [3–5]. PAACG begins with acute primary angle closure (APAC) attack and develops to optic neuropathy later [6, 7]. It was reported that high intraocular pressure (IOP) caused vascular insufficiency and ganglion cell apoptosis, leading to optic neuropathy and vision loss [6]. Recent reports also stated that PAACG-induced retinal nerve fiber layer (RNFL) thickness change was an evolving process between 2 and 16 weeks after the attack [8]. At present, the related examinations for glaucoma diagnosis and condition monitoring included visual acuity (VA), IOP, visual field (VF), and optical coherence tomography (SD-OCT). In addition, optical coherence tomography angiography (OCTA), as a new method, provides a more sensitive analysis on glaucoma diagnosis and mechanism at the vascular levels [9]. Previous studies reported the close relationship between these indicators [10], but the association between the PAACG attack time with the fundus condition is yet to be investigated.

The shallow anterior chamber depth is a common feature of glaucoma patients [11]. PAACG patients benefit a lot from the combined procedure of phacoemulsification and viscosgoniosynechialysis, which deal with cataract problem and expand the anterior chamber depth to reduce the patients' suddenly increased IOP at the same time [12]. This study aimed to investigate the structure and vessel changes of the RNFL of PAACG eyes from patients suffering from different attack times. Phacoemulsification and viscosgoniosynechialysis were performed after resolution of the acute episode in PAACG patients. In addition, VA, IOP, VF, and their correlations were investigated in our study. Our results indicated that shorter attack time resulted in better visual acuity, better VF, higher vascular density, thicker macula, and thicker RNFL.

2. Materials and Methods

2.1. Patients

Study subjects were recruited from patients with an episode of unilateral PAACG attack and cataracts at the Division of Glaucoma, Department of Ophthalmology, Zhongnan Hospital of Wuhan University, China, between January 2017 and April 2019. The study protocol was approved by the Institutional Review Board of Zhongnan Hospital of Wuhan University and adhered to the Declaration of Helsinki. The study protocol was registered on <http://www.ClinicalTrials.gov> (<https://clinicaltrials.gov/ct2/show/ChiCTR1800017849>). Written informed consent was obtained from each patient in this study prior to enrollment. The inclusion criteria for PAACG patients were as follows: (1) IOP > 21 mmHg with ≥ 2 following symptoms: visual blurring, ocular pain, headache, nausea, and vomiting; (2) the presence of any following signs on slit-lamp examination: conjunctival congestion, a mid-dilated unreactive pupil, corneal edema, and a shallow anterior chamber; (3) a closed angle observed by gonioscopy; (4) duration of PAACG attack < 120 hours; (5) IOP < 21 mmHg after medical therapy and surgery. The exclusion criteria were as follows: (1) bilateral attack; (2) secondary angle closure caused by phacomorphic angle closure, subluxed lens, and neovascularization; (3) a history of vitreoretinal or corneal surgery in the attack eye; (4) refraction > 5.0 D (sphere) and/or 3.0 D (cylinder); (5) incomplete clinical data, poor-quality OCT scans, the ratio of false negative or false positives in VF examination > 30%, or fixation loss of >20%; (6) Severe corneal edema after IOP control.

2.2. Examinations

Patients were divided into 2 groups according to their attack time: <1 day (group A) and ≥ 1 day (group B). All subjects underwent ophthalmic examinations including best-corrected visual acuity (BCVA, measured by Snellen chart), IOP measurement, slit-lamp examination, fundus examination, and gonioscopy. IOP was measured by iCare tonometry or Goldmann applanation tonometry. Gonioscopy was examined with a Goldmann 2-mirror lens at high magnification ($\times 16$). VF testing was performed using 24–2 pattern Swedish interactive threshold algorithm on the Humphrey Field Analyzer (Carl Zeiss, USA). VF indices included mean defect (MD) and pattern standard deviation (PSD). Normal VF was defined as MD and PSD within 95%

confidence intervals and a glaucoma hemifield test result "within normal limits." RNFL images were obtained using the stratus OCT 3000 (Carl Zeiss) with peripapillary RNFL thickness calculated automatically, including average, superior, nasal, inferior, and temporal quadrants. Only high-quality images with $3 \leq$ signal strength ≤ 10 were included. Macular thickness in scans with a 512×128 mm field and ganglion cell complex (GCC) thickness were also assessed with SD-OCT. Vessel perfusion density was measured within the cpVD in scans with a 6×6 mm field centered on the optic nerve head. Vessel density within the superficial retina layer was measured using the ZEISS CIRRUS™ OCT Angiography systemRTVue-XR Avanti.

2.3. Procedures

All patients underwent phacoemulsification and viscogoniosynechialysis. Two scheduled ophthalmic examinations were performed after IOP controlling: the first one before the procedure and the second one 1 month after the procedure. The patients with IOP > 21 mmHg 1 day after these procedures were further treated with IOP-lowering drugs, such as (1) beta-blockers, (2) carbonic anhydrase inhibitors, (3) pilocarpine, and (4) adrenergic agonists. Additional clinic visits and investigations were scheduled when required.

2.4. Statistical Analysis

All data were analyzed using the SPSS version 20.0. Independent-sample *T* test was used for the significance of differences between IOP, VA, OCT parameters, vessel flow density, and VF indices including MD and PSD. Pearson correlation was used to evaluate single correlation between average RNFL thickness, average GCC thickness, cpVD, and VF indices. The significance was set as a two-sided $p < 0.05$.

3. Results

3.1. Shorter Attack Time Indicated Better BCVA Improvement

A total of 26 eyes from 26 patients with unilateral PAACG were enrolled in this study. The characteristics of all the patients are listed in Table 1. Age ($p=0.092$) and gender ($p=0.855$) were comparable between groups. The mean duration of the PAACG attack was 10.0 ± 6.2 hours (range 2 to 19 h) and 79.3 ± 24.7 hours (range 28 to 120 h), respectively. The mean cpRNFL thickness, baseline vertical cup-to-disc ratio, mean GCC thickness, cpVD, and MD value from VF had no significant differences in the 2 groups (all, $p > 0.05$). The BCVA before and after surgery in both groups is listed in Table 2. The distributions of the initial and final logMAR VA were shown in Figure 1. Compared with the initial mean BCVA, the final mean BCVA was significantly improved in both groups. In addition, the group A (attack time < 1 day) had a better BCVA and greater improvement than the group B (attack time > 1 day).

Characteristics of patients with acute primary angle closure.

Characteristic (n = 26)	Group A (n = 13)	Group B (n = 13)	p*
Age	63.92 ± 8.06 (52~85)	69.46 ± 8.02 (57~82)	0.092
Sex (male/female)	6/7	7/6	0.855
Right eye/left eye	7/6	8/5	0.877
Duration of symptoms (hours)	10.0 ± 6.2 (2~19)	79.3 ± 24.7 (28~120)	0.001
Cataract surgery (yes/no)	13/0	13/0	1.000
IOP preoperative (mmHg)	49.54 ± 12.59 (29~78)	46.92 ± 9.68 (28~62)	0.558
IOP postoperative (mmHg)	12.77 ± 2.65 (9~17)	12.77 ± 3.85 (6~19)	0.834
Mean cpRNFL thickness	108.32 ± 10.25	100.46 ± 12.75	0.327
Vertical cup-to-disc ratio	0.56 ± 0.13	0.68 ± 0.23	0.087
Mean GCC thickness	85.32 ± 6.57	80.25 ± 8.27	0.076
cpVD	0.59 ± 0.06	0.55 ± 0.05	0.06

Characteristic (n = 26)	Group A (n = 13)	Group B (n = 13)	p*
MD	-2.57 ± 1.45	-3.85 ± 2.78	0.056

cpVD: circumpapillary retinal nerve fiber layer vessel density; MD: mean defect.

Comparison of postoperative characteristics between group A and group B.

	Group A (n = 13)	Group B (n = 13)	p*
BCVA (logMAR)			
Baseline	1.56 ± 0.89	0.97 ± 0.64	0.058
Final	0.25 ± 0.15	0.46 ± 0.51	0.167
Improvement	1.32 ± 0.84	0.50 ± 0.21	0.004
VF			
MD value	-3.65 ± 2.54	-16.05 ± 5.99	<0.001
PSD value	2.75 ± 1.10	9.43 ± 2.93	<0.001
Macular thickness (µm)			
Fovea area	255.00 ± 27.94	203.92 ± 59.73	0.010
Inner area	304.38 ± 23.41	254.23 ± 58.13	0.008
Outer area	269.92 ± 21.68	231.00 ± 32.39	0.001
GCC thickness (µm)	82.62 ± 8.76	65.23 ± 18.56	0.005
RNFL thickness (µm)			
Superior area	137.54 ± 12.23	103.92 ± 27.74	0.001
Temporal area	78.25 ± 11.93	75.15 ± 7.67	0.543
Inferior area	149.92 ± 21.73	100.38 ± 24.38	<0.001
Nasal area	68.08 ± 4.29	59.31 ± 14.67	0.057
Average area	105.08 ± 9.38	77.69 ± 20.23	<0.001
cpVD			
Central area	0.34 ± 0.03	0.29 ± 0.04	0.002
Inner area	0.56 ± 0.03	0.42 ± 0.06	<0.001
Outer area	0.56 ± 0.03	0.40 ± 0.08	<0.001
Full area	0.56 ± 0.02	0.41 ± 0.07	<0.001

BCVA: best-corrected visual acuity; logMAR = logarithm of the minimum angle of resolution. MD: mean defect; PSD: pattern standard deviation.

PHOTO (COLOR): Baseline and final logMAR BCVA of patients in this study. Dots below the line indicate improved visual acuity, and dots above the line indicate worse visual acuity.

3.2. Patients with Shorter Attack Time Had Better VF

As VF was measured 1 month after surgery, the MD and PSD values were significantly lower in patients from the group A than those from the group B (-3.65 ± 2.54 vs. -16.05 ± 5.99, p<0.001; -2.75 ± 1.10 vs. 9.43 ± 2.93, p<0.001; Table 2).

3.3. Patients with Shorter Attack Time Had Thicker Macula, GCC, and RNFL

The fovea and inner circle and outer circle thicknesses of macula were significantly different between groups 1 month after surgery (Table 2). One month after surgery, the macular fovea thickness in the group A was higher than that in group B (255.00 ± 27.94 µm vs. 203.92 ± 59.73 µm, p=0.010). The inner circle of macular thickness was higher in the group A (304.38 ± 23.41 µm vs. 254.23 ± 58.13, p=0.008). The outer circle of macular thickness was also higher in the group A (269.92 ± 21.68 µm vs. 231.00 ± 32.39 µm, p=0.001). These results indicated that the changes of macular thickness were correlated with attack time. Longer attack time indicated thinner macula. The average thickness of GCC was significantly higher in the group A than those in the group B 1 month after surgery (82.62 ± 8.76 µm vs. 65.23 ± 18.56 µm, p=0.005).

The superior, temporal, inferior, nasal, and average peripapillary RNFL thickness 1 month after surgery were listed in Table 2. Compared with group B, group A had a significantly thicker RNFL ($105.08 \pm 9.38 \mu\text{m}$ vs. $77.69 \pm 20.23 \mu\text{m}$, $p < 0.001$). Quadrants analyses showed higher superior ($137.54 \pm 12.23 \mu\text{m}$ vs. $103.92 \pm 27.74 \mu\text{m}$, $p = 0.001$) and inferior quadrants ($149.92 \pm 21.73 \mu\text{m}$ vs. $100.38 \pm 24.38 \mu\text{m}$, $p < 0.001$) in the group A but comparable thickness in the temporal and nasal quadrants ($p = 0.543$; $p = 0.057$).

3.4. Patients with Shorter Attack Time Had Higher cpVD

The OCTA-derived vessel flow densities from both groups are presented in Table 2. Eyes from the group A had significantly greater cpVD (circumpapillary RNFL vessel density) than those from the group B in the central area (0.34 ± 0.03 vs. 0.29 ± 0.04 ; $p = 0.002$), in the inner area (0.56 ± 0.03 vs. 0.42 ± 0.06 ; $p < 0.001$), in the outer area (0.56 ± 0.03 vs. 0.40 ± 0.08 ; $p < 0.001$), and in the full area (0.56 ± 0.02 vs. 0.41 ± 0.07 ; $p < 0.001$).

3.5. Relationship between VF Parameters, Average RNFL Thickness, and cpVD

The correlation coefficients between VF parameters, average RNFL thickness, and cpVD in both groups 1 month after surgery are summarized in Table 3. The significant correlations were found between all these parameters ($r > 0.4$, $p < 0.05$). In both groups, the association between the average RNFL thickness and cpVD, as well as MD values and PSD values from VF, was stronger ($R^2 = 0.58$, 0.60 , -0.54 , respectively, all, $p < 0.001$) than the association between the GCC thickness and cpVD, as well as MD values and PSD values ($R^2 = 0.37$, $p = 0.001$; $R^2 = 0.37$, $p = 0.001$; $R^2 = -0.27$, $p = 0.007$).

Pearson correlation for acute primary angle closure glaucoma eye.

Variable	cpVD	VF MD	VF PSD
Mean RNFL thickness	$r = 0.761$, $p < 0.001$	$r = 0.772$, $p < 0.001$	$r = -0.736$, $p < 0.001$
Mean GCC thickness	$r = 0.605$, $p = 0.001$	$r = 0.605$, $p = 0.001$	$r = -0.516$, $p = 0.007$
cpVD		$r = 0.832$, $p < 0.001$	$r = -0.803$, $p < 0.001$

3.6. Patients with Shorter Attack Time Had Better Retinal Function

The fundus conditions of 2 representative patients from both groups are shown in Figure 2. The visual function of the patient from the group A was better than that from the group B.

PHOTO (COLOR): Representative retinal damage images of 2 patients from different groups. (a) cpVD: OCT angiography image of a 6×6 mm section, centered on the optic disc head. Group A showed a better blood flow density than group B (yellow arrowheads). (b) ONH: RNFL thickness analysis examined by OCT. Group B was detected a more apparent RNFL defect in the superior region than group A (blue arrowheads). (c) GCC: ganglion cell complex thickness. An apparent GCC defect was detected in the inferotemporal area (red arrowheads). (d) MD: VF pattern deviation plot. Superior visual field loss was detected in group B patient.

4. Discussion

In our study, we used the diagnostic ability of BCVA and structural and vascular changes from macula and RNFL to detect retinal changes in PAACG patients. Most patients had good visual outcomes after surgery, and patients with shorter attack time (less than 1 day) got better BCVA. Patients with shorter attack time had thicker macula, GCC, and RNFL. Moreover, the retinal vessel densities in the peripapillary areas were significantly greater in the patients with shorter attack time.

SD-OCT and OCT-A were recently developed to observe the retinal function in glaucoma patients in both the foveal and disc areas, in an accurate, convenient, and noninvasive way [13, 14]. Reduced blood flow in the choroid, retina, and optic nerve head in glaucoma patients was proved using OCT angiography [15–18]. Our results showed that the RNFL thickness was higher in PAACG patients at the early stage of relief. This

thickening is commonly due to retinal edema. The results of our studies also demonstrated that the retinal thickness of all PAACG patients slowly decreases with the passage of time, which was consistent with the findings of Rao et al. [19] and Wang et al. [20]. Our results suggested that the speed and extent of retinal function loss were closely proportional to the attack time. These results might be related to 2 classic theories: a mechanical theory and a vascular theory [21, 22]. The mechanical theory noted that increased IOP led to elongation, stretching, and collapse of the laminar beams and retinal ganglion cell (RGC) axons damage [23]. According to the vascular theory, insufficient blood supply on account of either increased IOP or other risk factors reducing ocular blood flow may cause glaucomatous optic neuropathy (GON) [24]. Hayreh et al. also noted that the retinal circulation was not easily damaged at 30 mmHg IOP, unless IOP reached 40–70 mmHg. When IOP increased to 70 mmHg, the choroid capillaries in disc area almost disappeared, resulting in slow retinal circulation. These theories and phenomena may explain why prolonged periods of high IOP have significant effects on the fundus of PAACG patients.

In our studies, the thickness of the retina was significantly reduced on the superior and inferior areas, while the changes of nasal side and temporal side in the RNFL were lighter. These results were consistent with the findings of Igarashi et al. [9], who reported that the superior and inferior areas of the optic disc were more affected by glaucoma, which was the pathological characteristic of glaucomatous optic neuropathy [18].

In this study, we also evaluated the relationship between VF and fundus changes. There was a strong correlation between the VF loss and cpVD (all, $p < 0.001$), suggesting that reduced vessel flow density was associated with more advanced stages of glaucomatous VF damage. Moreover, we also found that changes in RNFL and RGC thickness often occurred later than changes in VF and blood flow density [25, 26]. There are several possible explanations for the independent association of vascular density and the MD of VF: first, it may be due to the reduced and dysfunctional RGCs (i.e., preapoptosis). Blood flow is therefore lower in vascular density and less sensitive to VF. Since these dysfunctional RGCs did not shrink, and the reduction in RNFL thickness and marginal area might not be discovered [10]. In addition, histological studies showed that there was only moderate correlation between RNFL thinning and RGC loss, indicating that RNFL thinning did not fully reflect the RGC function [5, 27]. Thus, a stronger correlation between vessel density and VF impairment indicated that vessel flow density was a better reflection of RGC function than structural loss [28]. In other words, these 2 indicators may play major roles in detecting early stage of glaucoma.

Our research had several limitations that need to be considered. All patients were selected from a single institution; therefore, a recommendation bias might exist. The number of patients enrolled in this study was limited, and this study lacked long-term follow-up of this series of data on the fundus, such as macular/GCC/RNFL thickness, VF, and cpVD, which might result in further fundus changes to the PAACG patients. However, from our current research data, it was enough to show that for PAACG patients, the attack time and retinal changes were clearly correlated.

According to our results, cpVD had more obvious changes than RNFL thickness, macular thickness, and GCC thickness. We considered that OCT-A vessel indicators may be more sensitive than SD-OCT at the early stage of PAACG. Therefore, OCT-A is a promising technology for clinical monitoring of vascular changes in glaucoma. Its application may help the ophthalmologist further understand the pathophysiology of glaucoma and its underlying vascular mechanisms.

In summary, longer attack time indicated longer periods of high IOP and perfusion damage to the retinal vessels. It not only caused short-term damage to the fundus but also played a negative role in subsequent

fundus change process. Therefore, it is importantly necessary to control the IOP in time once the PAACG patients had the relevant symptoms.

Data Availability

The clinical data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

References

- 1 Tan G. S. W., Hoh S. T., Husain R., *Visual acuity after acute primary angle closure and considerations for primary lens extraction, British Journal of Ophthalmology, 2006, 90, 1, 14, 16, 10.1136/bjo.2005.077115, 2-s2.0-29744436670*
- 2 Ch'ng T. W., Mosavi S. A., Noor Azimah A. A., Azlan N. Z., Azhany Y., Liza-Sharmini A. T., *Monsoon and primary acute angle closure in Malaysia, Medical Journal of Malaysia, 2013, 68, 5, 410, 414*
- 3 Lowe R. F., *Clinical types of primary angle closure glaucoma, Australian and New Zealand Journal of Ophthalmology, 1988, 16, 3, 245, 250, 10.1111/j.1442-9071.1988.tb01216.x, 2-s2.0-0024271316*
- 4 Amerasinghe N., Aung T., *Angle-closure: risk factors, diagnosis and treatment, Progress in Brain Research, 2008, 173, 31, 45, 10.1016/s0079-6123(08)01104-7, 2-s2.0-53049093164*
- 5 Tzamalís A., Pham D.-T., Wirbelauer C., *Comparison of slit lamp-adapted optical coherence tomography features of fellow eyes of acute primary angle closure and eyes with open angle glaucoma, Japanese Journal of Ophthalmology, 2014, 58, 3, 252, 260, 10.1007/s10384-013-0301-5, 2-s2.0-84900799687*
- 6 Foster P. J., Buhrmann R., Quigley H. A., Johnson G. J., *The definition and classification of glaucoma in prevalence surveys, British Journal of Ophthalmology, 2002, 86, 2, 238, 242, 10.1136/bjo.86.2.238, 2-s2.0-0036158752*
- 7 Muto T., Nishimura T., Sakamoto M., Inomata T., Machida S., *Identification of eyes at risk of acute primary angle-closure in elderly Japanese patients, Clinical Ophthalmology, 2019, 13, 859, 868, 10.2147/opth.s190942*
- 8 Lee J. W. Y., Lai J. S. M., Yick D. W. F., Yuen C. Y. F., *Prospective study on retinal nerve fibre layer changes after an acute episode of phacomorphic angle closure, International Ophthalmology, 2012, 32, 6, 577, 582, 10.1007/s10792-012-9614-5, 2-s2.0-84872724302*
- 9 Igarashi R., Ochiai S., Sakaue Y., *Optical coherence tomography angiography of the peripapillary capillaries in primary open-angle and normal-tension glaucoma, PLoS One, 2017, 12, 9, e0184301, 10.1371/journal.pone.0184301, 2-s2.0-85029891521*
- Yarmohammadi A., Zangwill L. M., Diniz-Filho A., *Relationship between optical coherence tomography angiography vessel density and severity of visual field loss in glaucoma, Ophthalmology, 2016, 123, 12, 2498, 2508, 10.1016/j.opthta.2016.08.041, 2-s2.0-84992316836*
- Salmon J. F., *Predisposing factors for chronic angle-closure glaucoma, Progress in Retinal and Eye Research, 1998, 18, 1, 121, 132, 10.1016/s1350-9462(98)00007-x, 2-s2.0-0032889609*

- Zhang Z., Chen M., Yin J., Yao K., *Microcoaxial phacoemulsification combined with viscogoniosynechialysis for patients with refractory acute angle-closure glaucoma*, *Journal of Glaucoma*, 2014, 23, 2, 119, 124, 10.1097/ijg.0b013e318268518d, 2-s2.0-84894072051
- Zhu L., Zong Y., Yu J., *Reduced retinal vessel density in primary angle closure glaucoma*, *Journal of Glaucoma*, 2018, 27, 4, 322, 327, 10.1097/ijg.0000000000000900, 2-s2.0-85045047816
- Ye C., Yu M., Leung C. K.-S., *Impact of segmentation errors and retinal blood vessels on retinal nerve fibre layer measurements using spectral-domain optical coherence tomography*, *Acta Ophthalmologica*, 2016, 94, 3, e211, e219, 10.1111/aos.12762, 2-s2.0-84934775141
- Hitchings R. A., Spaeth G. L., *Fluorescein angiography in chronic simple and low-tension glaucoma*, *British Journal of Ophthalmology*, 1997, 61, 126, 132, 10.1136/bjo.61.2.126, 2-s2.0-0017642024
- Tanaka Y., *Color-fluorescein relationship in glaucomatous optic nerve damage*, *Japanese Journal of Ophthalmology*, 1995, 39, 2, 180, 186
- Schwartz B., Rieser J. C., Fishbein S. L., *Fluorescein angiographic defects of the optic disc in glaucoma*, *Archives of Ophthalmology*, 1977, 95, 11, 1961, 1974, 10.1001/archophth.1977.04450110055002, 2-s2.0-0017551138
- Moghim S., SafiZadeh M., Fard M. A., *Changes in optic nerve head vessel density after acute primary angle closure episode*, *Investigative Ophthalmology & Visual Science*, 2019, 60, 2, 552, 10.1167/iovs.18-25915, 2-s2.0-85061146340
- Rao H. L., Pradhan Z. S., Weinreb R. N., *Vessel density and structural measurements of optical coherence tomography in primary angle closure and primary angle closure glaucoma*, *American Journal of Ophthalmology*, 2017, 177, 106, 115, 10.1016/j.ajo.2017.02.020, 2-s2.0-85015445120
- Wang X., Jiang C., Kong X., Yu X., Sun X., *Peripapillary retinal vessel density in eyes with acute primary angle closure: an optical coherence tomography angiography study*, *Graefe's Archive for Clinical and Experimental Ophthalmology*, 2017, 255, 5, 1013, 1018, 10.1007/s00417-017-3593-1, 2-s2.0-85011607390
- Hayreh S. S., *Optic disc changes in glaucoma*, *British Journal of Ophthalmology*, 1972, 56, 3, 175, 185, 10.1136/bjo.56.3.175, 2-s2.0-0015307533
- Fechtner R. D., Weinreb R. N., *Mechanisms of optic nerve damage in primary open angle glaucoma*, *Survey of Ophthalmology*, 1994, 39, 1, 23, 42, 10.1016/s0039-6257(05)80042-6, 2-s2.0-0028038213
- Yan D. B., Coloma F. M., Metheerairut A., Trope G. E., Heathcote J. G., Ethier C. R., *Deformation of the lamina cribrosa by elevated intraocular pressure*, *British Journal of Ophthalmology*, 1994, 78, 8, 643, 648, 10.1136/bjo.78.8.643, 2-s2.0-0027969262
- Flammer J., *The vascular concept of glaucoma*, *Survey of Ophthalmology*, 1994, 38, S3, S6, 10.1016/0039-6257(94)90041-8, 2-s2.0-0028433377
- Wang X., Jiang C., Ko T., *Correlation between optic disc perfusion and glaucomatous severity in patients with open-angle glaucoma: an optical coherence tomography angiography study*, *Graefe's Archive for Clinical and Experimental Ophthalmology*, 2015, 253, 9, 1557, 1564, 10.1007/s00417-015-3095-y, 2-s2.0-84940436047

Yarmohammadi A., Zangwill L. M., Diniz-Filho A., *Optical coherence tomography angiography vessel density in healthy, glaucoma suspect, and glaucoma eyes, Investigative Ophthalmology & Visual Science, 2016, 57, 9, OCT451, OCT459, 10.1167/iovs.15-18944, 2-s2.0-84978531727*

Chauhan B. C., Stevens K. T., Levesque J. M., *Longitudinal in vivo imaging of retinal ganglion cells and retinal thickness changes following optic nerve injury in mice, PLoS One, 2012, 7, 6, e400352, 10.1371/journal.pone.0040352, 2-s2.0-84863109512*

Sehi M., Goharian I., Konduru R., *Retinal blood flow in glaucomatous eyes with single-hemifield damage, Ophthalmology, 2014, 121, 3, 750, 758, 10.1016/j.ophtha.2013.10.022, 2-s2.0-84896705662*

~~~~~

By Xiaolu Zhu; Wen Zeng; Shengyu Wu; Xiaomin Chen; Tian Zheng and Min Ke

---

Copyright of Journal of Ophthalmology is the property of Hindawi Limited and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.

**Record: 1**

**Title:** Self-Monitoring Symptoms in Glaucoma: A Feasibility Study of a Web-Based Diary Tool.

**Authors:** McDonald L; Division of Optometry and Visual Sciences, School of Health Sciences, City, University of London, London EC1V 0HB, UK.  
Glen FC; Division of Optometry and Visual Sciences, School of Health Sciences, City, University of London, London EC1V 0HB, UK.  
Taylor DJ; Division of Optometry and Visual Sciences, School of Health Sciences, City, University of London, London EC1V 0HB, UK.  
Crabb DP; Division of Optometry and Visual Sciences, School of Health Sciences, City, University of London, London EC1V 0HB, UK.

**Source:** Journal of ophthalmology [J Ophthalmol] 2017; Vol. 2017, pp. 8452840.  
*Date of Electronic Publication:* 2017 May 04.

**Publication Type:** Journal Article

**Language:** English

**Journal Info:** *Publisher:* Hindawi Publishing Corporation *Country of Publication:* United States *NLM ID:* 101524199 *Publication Model:* Print-Electronic *Cited Medium:* Print *ISSN:* 2090-004X (Print) *Linking ISSN:* 2090004X *NLM ISO Abbreviation:* J Ophthalmol *Subsets:* PubMed not MEDLINE

**Imprint Name(s):** *Original Publication:* New York, NY : Hindawi Publishing Corporation

**Abstract:** Purpose . Glaucoma patients annually spend only a few hours in an eye clinic but spend more than 5000 waking hours engaged in everything else. We propose that patients could self-monitor changes in visual symptoms providing valuable between clinic information; we test the hypothesis that this is feasible using a web-based diary tool. Methods . Ten glaucoma patients with a range of visual field loss took part in an eight-week pilot study. After completing a series of baseline tests, volunteers were prompted to monitor symptoms every three days and complete a diary about their vision during daily life using a bespoke web-based diary tool. Response to an end of a study questionnaire about the usefulness of the exercise was a main outcome measure. Results . Eight of the 10 patients rated the monitoring scheme to be "valuable" or "very valuable." Completion rate to items was excellent (96%). Themes from a qualitative synthesis of the diary entries related to behavioural aspects of glaucoma. One patient concluded that a constant focus on monitoring symptoms led to negative feelings. Conclusions . A web-based diary tool for monitoring self-reported glaucoma symptoms is practically feasible. The tool must be carefully designed to ensure participants are benefitting, and it is not increasing anxiety.

**References:** Comput Methods Programs Biomed. 2002 Jul;69(1):75-86. (PMID: 12088595)  
Ann Med. 2001 Jul;33(5):337-43. (PMID: 11491192)  
Int J Behav Nutr Phys Act. 2014 Jul 25;11:97. (PMID: 25059981)

Diabet Med. 2005 Oct;22(10):1372-8. (PMID: 16176199)  
Br J Ophthalmol. 2015 Mar;99(3):313-7. (PMID: 25336582)  
Invest Ophthalmol Vis Sci. 2011 Sep 01;52(9):7007-11. (PMID: 21810985)  
BMC Ophthalmol. 2015 Oct 06;15:128. (PMID: 26445483)  
Ann Intern Med. 2012 Mar 20;156(6):416-24. (PMID: 22431674)  
JAMA Ophthalmol. 2014 Jul;132(7):845-50. (PMID: 24831037)  
Graefes Arch Clin Exp Ophthalmol. 2005 Mar;243(3):210-6. (PMID: 15806374)  
N Engl J Med. 2012 Jul 5;367(1):1-3. (PMID: 22716935)  
Am J Geriatr Psychiatry. 2005 Mar;13(3):180-7. (PMID: 15728748)  
Ophthalmology. 2008 Aug;115(8):1320-7, 1327.e1-3. (PMID: 18321582)  
BMC Health Serv Res. 2013 Jul 03;13:251. (PMID: 23819721)  
Arch Ophthalmol. 2004 Jan;122(1):84-8. (PMID: 14718300)  
Psychosom Med. 2003 Nov-Dec;65(6):963-70. (PMID: 14645773)  
Psychol Bull. 2002 Jul;128(4):638-62. (PMID: 12081086)  
Arch Ophthalmol. 1997 Nov;115(11):1447-55. (PMID: 9366678)  
Trials. 2015 Nov 11;16:517. (PMID: 26559142)  
Nat Med. 2014 Sep;20(9):1074-8. (PMID: 25150497)  
Am J Ophthalmol. 2006 Jan;141(1):24-30. (PMID: 16386972)  
Gen Hosp Psychiatry. 2008 May-Jun;30(3):208-25. (PMID: 18433653)  
Ann Intern Med. 1997 Dec 15;127(12):1097-102. (PMID: 9412313)  
Ophthalmology. 2009 Dec;116(12):2286-93. (PMID: 19815286)  
J Glaucoma. 2003 Apr;12(2):139-50. (PMID: 12671469)  
Ophthalmology. 2016 Aug;123(8):1675-84. (PMID: 27289178)  
BMJ. 2008 Jun 28;336(7659):1464-5. (PMID: 18583674)  
Ophthalmology. 2009 Dec;116(12):2277-85.e1-3. (PMID: 19744715)  
Sociol Health Illn. 2007 Nov;29(7):1059-74. (PMID: 18092983)  
BMC Ophthalmol. 2011 Aug 04;11:19. (PMID: 21816060)

**Entry Date(s):** *Date Created:* 20170527 *Latest Revision:* 20200930

**Update Code:** 20210210

**PubMed Central ID:** PMC5435987

**DOI:** 10.1155/2017/8452840

**PMID:** 28546876

**Database:** MEDLINE with Full Text