

Your Name:

Isomer Relationships, Conformational Stability, and Reaction Selectivity

Instructions:

Read the instructions carefully and answer the following questions in the space provided. Then attach the PDF file on Discussion Board for peer discussion and peer review.

1. In a few sentences define each of the three main classes of isomers below. Configurational Isomers can be further classified as either enantiomers or diastereomers and should also be discussed. Your descriptions should enable your peers to quickly and easily compare and contrast the differences between the different classes of isomers.
 - A. Constitutional Isomers (or Structural Isomers)
 - B. Conformational Isomers
 - C. Configurational Isomers (or Stereoisomers)
 - I. Enantiomers
 - II. Diastereomers

- A. Constitutional Isomers - They have the same molecular formula but a different bonding arrangement among atoms

B. Conformational Isomers - Same bond connectivity, same configuration, different spatial arrangement - produced by rotation






C. Configurational Isomers - Same connectivity, different arrangement in space.

(i) Enantiomers - They are mirror images of one another


(ii) Diastereomers - They are not mirror images of one another

-
2. Analyze each pair of compounds below. Identify which class of isomers each pair of structures represent. Your choices are: *(i) constitutional, (ii) conformational, (iii) enantiomers, or (iv) diastereomers or (v) identical (non-isomers)*. Only one label for each pair should be used. For each case show work (drawings, (R)/(S), rotation of bonds, chair-flips, etc) which led you to each answer for the pairs below. [During your peer review, any differences of answers between you and your peers can be discussed and corrected as needed.]


Note: Various projections may be used when drawing chemical structures including bond-line structures (using proper dash/wedge bonds), Newman projections, chair structures (for cyclohexane derivatives), Haworth projections, and Fischer projections. These projections just represent different vantage points of visualizing a chemical structure and can be interconverted from one projection to another. Each different projection has been labeled for you below. These are all utilized throughout the text book.




- This represents constitutional isomers due to the different connectivity





Redrawing:



- They are the same
Therefore, they are:
identical (non-isomers)



- This represents enantiomers because they are mirror images of one another.




- This represents diastereomers because they have same connectivity and formula, and they are NOT mirror images of one another


3. Analyze the structures for **Compound A** and **Compound B** shown below. Answer and discuss each question (a) and (b), respectively.

- What type of isomers do these represent?
- Which compound is more stable? (Hint: You will need to draw the chair structures for each compound (4 total chairs, two for each compound) and compare/contrast/analyze the various 1,3-diaxial interactions in order to justify this answer. Show your work clearly.)

Compound A





Compound B



(a) Type of isomers


- Compounds A and B represent constitutional isomers as they have a different bonding arrangement & same formula.

(b) Compound B is more stable



- B is more stable since the first structure does not have 1,3-diaxial interactions


4. Chiral catalysts can be designed to favor the formation of one enantiomer over another in reactions where a new chiral center is formed. The example shown below was recently reported in the chemical literature (*J. Am. Chem. Soc.* **2012**, *134*, 8054–8057) where a new type of chiral copper (Cu) catalyst was developed. When the Cu^{I} catalyst was used, the (S)-enantiomer of the product predominated, and the (R)-enantiomer of the product was selectively formed when Cu^{II} catalyst was used instead.




Answer part (a) and (b) below. You aren't required to understand the chemistry to answer correctly and discuss the experimental results.

- a. The product has been shown on the right of the reaction scheme above but the absolute stereochemistry of the product has been excluded. Draw the **major enantiomer** formed when each form of the catalyst is used. You need to be sure to show the stereochemistry clearly using appropriate dash/wedge bonds at the stereocenter and indicate which catalyst (Cu^{I} or Cu^{II}) was used for each of the two individual enantiomers you draw.

(i) When Cu^{I} is used, the (S)-enantiomer is formed. This is:



(ii) (R)-enantiomer is formed when Cu^{II} is used:



- b. The chart below reports the data when the (S)-selective catalyst was used in various solvents. The enantiomeric excess (reported as % ee) is shown for each case.

Solvent	% ee of (S)-product	% Yield of total product	% S	% R
Toluene	24	55	62	38
THF	48	33	74	26
Acetonitrile	72	55	86	14
Chloroform	30	40	65	35
Dichloromethane	46	44	73	27
Hexane	51	30	75.5	24.5

Complete the table above by calculating the % (S) and % (R) formed when the (S)-selective reaction was repeated in the various solvents. How are these values calculated? Illustrate this by showing a complete calculation for the toluene case. For the others, you can simply just write in the values. If your values do not match your peers', you will need to discuss this and correct as needed. Also, discuss which solvent gives the optimal results in terms of both enantioselectivity and overall total product yield. What experimental data led you to make this conclusion? Where any other solvents comparable? What solvent gave the worst results?

Toluene case: $S + R = 100$. Therefore, $R = S - 24$

$$\& \& R + 24 = S \quad | \&$$

Substituting $(S - 24)$ in the eqn $S + R = 100$:

$$S + (S - 24) = 100$$

$$S = 62 \text{ and } R = 38$$

- The solvent that gives optimal results is Acetonitrile because it has the highest value of enantioselectivity.

- Toluene has a high yield even though it has the lowest $\& \& \% S$. On the

The solvent that gave the worst results is chloroform due to its relatively low yield and $\& \& \% S$.