

Major Depressive Disorder

Diagnostic Criteria

- A. Five (or more) of the following symptoms have been present during the same 2-week period and represent a change from previous functioning; at least one of the symptoms is either (1) depressed mood or (2) loss of interest or pleasure.

Note: Do not include symptoms that are clearly attributable to another medical condition.

1. Depressed mood most of the day, nearly every day, as indicated by either subjective report (e.g., feels sad, empty, hopeless) or observation made by others (e.g., appears tearful). (**Note:** In children and adolescents, can be irritable mood.)
2. Markedly diminished interest or pleasure in all, or almost all, activities most of the day, nearly every day (as indicated by either subjective account or observation).

3. Significant weight loss when not dieting or weight gain (e.g., a change of more than 5% of body weight in a month), or decrease or increase in appetite nearly every day. (**Note:** In children, consider failure to make expected weight gain.)
 4. Insomnia or hypersomnia nearly every day.
 5. Psychomotor agitation or retardation nearly every day (observable by others, not merely subjective feelings of restlessness or being slowed down).
 6. Fatigue or loss of energy nearly every day.
 7. Feelings of worthlessness or excessive or inappropriate guilt (which may be delusional) nearly every day (not merely self-reproach or guilt about being sick).
 8. Diminished ability to think or concentrate, or indecisiveness, nearly every day (either by subjective account or as observed by others).
 9. Recurrent thoughts of death (not just fear of dying), recurrent suicidal ideation without a specific plan, or a suicide attempt or a specific plan for committing suicide.
- B. The symptoms cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.
- C. The episode is not attributable to the physiological effects of a substance or another medical condition.

Note: Criteria A–C represent a major depressive episode.

Note: Responses to a significant loss (e.g., bereavement, financial ruin, losses from a natural disaster, a serious medical illness or disability) may include the feelings of intense sadness, rumination about the loss, insomnia, poor appetite, and weight loss noted in Criterion A, which may resemble a depressive episode. Although such symptoms may be understandable or considered appropriate to the loss, the presence of a major depressive episode in addition to the normal response to a significant loss should also be carefully considered. This decision inevitably requires the exercise of clinical judgment based on the individual's history and the cultural norms for the expression of distress in the context of loss.¹

- D. The occurrence of the major depressive episode is not better explained by schizoaffective disorder, schizophrenia, schizophreniform disorder, delusional disorder, or other specified and unspecified schizophrenia spectrum and other psychotic disorders.
- E. There has never been a manic episode or a hypomanic episode.

Note: This exclusion does not apply if all of the manic-like or hypomanic-like episodes are substance-induced or are attributable to the physiological effects of another medical condition.

¹In distinguishing grief from a major depressive episode (MDE), it is useful to consider that in grief the predominant affect is feelings of emptiness and loss, while in an MDE it is persistent depressed mood and the inability to anticipate happiness or pleasure. The dysphoria in grief is likely to decrease in intensity over days to weeks and occurs in waves, the so-called pangs of grief. These waves tend to be associated with thoughts or reminders of the deceased. The depressed mood of an MDE is more persistent and not tied to specific thoughts or preoccupations. The pain of grief may be accompanied by positive emotions and humor that are uncharacteristic of the pervasive unhappiness and misery characteristic of an MDE. The thought content associated with grief generally features a preoccupation with thoughts and memories of the deceased, rather than the self-critical or pessimistic ruminations seen in an MDE. In grief, self-esteem is generally preserved, whereas in an MDE feelings of worthlessness and self-loathing are common. If self-derogatory ideation is present in grief, it typically involves perceived failings vis-à-vis the deceased (e.g., not visiting frequently enough, not telling the deceased how much he or she was loved). If a bereaved individual thinks about death and dying, such thoughts are generally focused on the deceased and possibly about "joining" the deceased, whereas in an MDE such thoughts are focused on ending one's own life because of feeling worthless, undeserving of life, or unable to cope with the pain of depression.

Coding and Recording Procedures

The diagnostic code for major depressive disorder is based on whether this is a single or recurrent episode, current severity, presence of psychotic features, and remission status. Current severity and psychotic features are only indicated if full criteria are currently met for a major depressive episode. Remission specifiers are only indicated if the full criteria are not currently met for a major depressive episode. Codes are as follows:

Severity/course specifier	Single episode	Recurrent episode*
Mild (p. 188)	296.21 (F32.0)	296.31 (F33.0)
Moderate (p. 188)	296.22 (F32.1)	296.32 (F33.1)
Severe (p. 188)	296.23 (F32.2)	296.33 (F33.2)
With psychotic features** (p. 186)	296.24 (F32.3)	296.34 (F33.3)
In partial remission (p. 188)	296.25 (F32.4)	296.35 (F33.41)
In full remission (p. 188)	296.26 (F32.5)	296.36 (F33.42)
Unspecified	296.20 (F32.9)	296.30 (F33.9)

*For an episode to be considered recurrent, there must be an interval of at least 2 consecutive months between separate episodes in which criteria are not met for a major depressive episode. The definitions of specifiers are found on the indicated pages.

**If psychotic features are present, code the "with psychotic features" specifier irrespective of episode severity.

In recording the name of a diagnosis, terms should be listed in the following order: major depressive disorder, single or recurrent episode, severity/psychotic/remission specifiers, followed by as many of the following specifiers without codes that apply to the current episode.

Specify:

- With anxious distress (p. 184)
- With mixed features (pp. 184–185)
- With melancholic features (p. 185)
- With atypical features (pp. 185–186)
- With mood-congruent psychotic features (p. 186)
- With mood-incongruent psychotic features (p. 186)
- With catatonia (p. 186). **Coding note:** Use additional code 293.89 (F06.1).
- With peripartum onset (pp. 186–187)
- With seasonal pattern (recurrent episode only) (pp. 187–188)

Diagnostic Features

The criterion symptoms for major depressive disorder must be present nearly every day to be considered present, with the exception of weight change and suicidal ideation. Depressed mood must be present for most of the day, in addition to being present nearly every day. Often insomnia or fatigue is the presenting complaint, and failure to probe for accompanying depressive symptoms will result in underdiagnosis. Sadness may be denied at first but may be elicited through interview or inferred from facial expression and demeanor. With individuals who focus on a somatic complaint, clinicians should determine whether the distress from that complaint is associated with specific depressive symptoms. Fatigue and sleep disturbance are present in a high proportion of cases; psychomotor disturbances are much less common but are indicative of greater overall severity, as is the presence of delusional or near-delusional guilt.

The essential feature of a major depressive episode is a period of at least 2 weeks during which there is either depressed mood or the loss of interest or pleasure in nearly all activities (Criterion A). In children and adolescents, the mood may be irritable rather than sad. The individual must also experience at least four additional symptoms drawn from a list that includes changes in appetite or weight, sleep, and psychomotor activity; decreased energy; feelings of worthlessness or guilt; difficulty thinking, concentrating, or making decisions; or recurrent thoughts of death or suicidal ideation or suicide plans or attempts. To count toward a major depressive episode, a symptom must either be newly present or must have clearly worsened compared with the person's pre-episode status. The symptoms must persist for most of the day, nearly every day, for at least 2 consecutive weeks. The episode must be accompanied by clinically significant distress or impairment in social, occupational, or other important areas of functioning. For some individuals with milder episodes, functioning may appear to be normal but requires markedly increased effort.

The mood in a major depressive episode is often described by the person as depressed, sad, hopeless, discouraged, or "down in the dumps" (Criterion A1). In some cases, sadness may be denied at first but may subsequently be elicited by interview (e.g., by pointing out that the individual looks as if he or she is about to cry). In some individuals who complain of feeling "blah," having no feelings, or feeling anxious, the presence of a depressed mood can be inferred from the person's facial expression and demeanor. Some individuals emphasize somatic complaints (e.g., bodily aches and pains) rather than reporting feelings of sadness. Many individuals report or exhibit increased irritability (e.g., persistent anger, a tendency to respond to events with angry outbursts or blaming others, an exaggerated sense of frustration over minor matters). In children and adolescents, an irritable or cranky mood may develop rather than a sad or dejected mood. This presentation should be differentiated from a pattern of irritability when frustrated.

Loss of interest or pleasure is nearly always present, at least to some degree. Individuals may report feeling less interested in hobbies, "not caring anymore," or not feeling any enjoyment in activities that were previously considered pleasurable (Criterion A2). Family members often notice social withdrawal or neglect of pleasurable avocations (e.g., a formerly avid golfer no longer plays, a child who used to enjoy soccer finds excuses not to practice). In some individuals, there is a significant reduction from previous levels of sexual interest or desire.

Appetite change may involve either a reduction or increase. Some depressed individuals report that they have to force themselves to eat. Others may eat more and may crave specific foods (e.g., sweets or other carbohydrates). When appetite changes are severe (in either direction), there may be a significant loss or gain in weight, or, in children, a failure to make expected weight gains may be noted (Criterion A3).

Sleep disturbance may take the form of either difficulty sleeping or sleeping excessively (Criterion A4). When insomnia is present, it typically takes the form of middle insomnia (i.e., waking up during the night and then having difficulty returning to sleep) or terminal insomnia (i.e., waking too early and being unable to return to sleep). Initial insomnia (i.e., difficulty falling asleep) may also occur. Individuals who present with oversleeping (hypersomnia) may experience prolonged sleep episodes at night or increased daytime sleep. Sometimes the reason that the individual seeks treatment is for the disturbed sleep.

Psychomotor changes include agitation (e.g., the inability to sit still, pacing, hand-wringing; or pulling or rubbing of the skin, clothing, or other objects) or retardation (e.g., slowed speech, thinking, and body movements; increased pauses before answering; speech that is decreased in volume, inflection, amount, or variety of content, or muteness) (Criterion A5). The psychomotor agitation or retardation must be severe enough to be observable by others and not represent merely subjective feelings.

Decreased energy, tiredness, and fatigue are common (Criterion A6). A person may report sustained fatigue without physical exertion. Even the smallest tasks seem to require