




James Earl Jones

WHAT IS INVOKANA® (canagliflozin)?

INVOKANA® is a prescription medicine used along with diet and exercise to lower blood sugar in adults with type 2 diabetes. INVOKANA® is not for people with type 1 diabetes or with diabetic ketoacidosis (increased ketones in blood or urine). It is not known if INVOKANA® is safe and effective in children under 18 years of age.

IMPORTANT SAFETY INFORMATION

INVOKANA® can cause important side effects, including:

- **Dehydration.** INVOKANA® can cause some people to become dehydrated (the loss of too much body water), which may cause you to feel dizzy, faint, lightheaded, or weak, especially when you stand up (orthostatic hypotension). You may be at higher risk of dehydration if you have low blood pressure, take medicines to lower your blood pressure (including diuretics [water pills]), are on a low sodium (salt) diet, have kidney problems, or are 65 years of age or older
- **Vaginal yeast infection.** Women who take INVOKANA® may get vaginal yeast infections. Symptoms include: vaginal odor, white or yellowish vaginal discharge (discharge may be lumpy or look like cottage cheese), or vaginal itching
- **Yeast infection of the penis (balanitis or balanoposthitis).** Men who take INVOKANA® may get a yeast infection of the skin around the penis. Symptoms include: redness, itching, or swelling of the penis; rash of the penis; foul-smelling discharge from the penis; or pain in the skin around penis

Talk to your doctor about what to do if you get symptoms of a yeast infection of the vagina or penis.

Do not take INVOKANA® if you:

- are allergic to canagliflozin or any of the ingredients in INVOKANA®. Symptoms of allergic reaction may include: rash; raised red patches on your skin (hives); or swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing
- have severe kidney problems or are on dialysis

Before you take INVOKANA®, tell your doctor if you have kidney problems; liver problems; history of urinary tract infections or problems with urination; are on a low sodium (salt) diet; are going to have surgery; are eating less due to illness, surgery, or change in diet; pancreas problems; drink alcohol very often (or drink a lot of alcohol in short-term); ever had an allergic reaction to INVOKANA®; or have other medical conditions.

Tell your doctor if you are or plan to become pregnant, are breastfeeding, or plan to breastfeed. INVOKANA® may harm your unborn baby. If you become pregnant while taking INVOKANA®, tell your doctor right away. INVOKANA® may pass into your breast milk and may harm your baby. Do not breastfeed while taking INVOKANA®.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Especially tell your doctor if you take diuretics (water pills), rifampin (used to treat or prevent tuberculosis), phenytoin or phenobarbital (used to control seizures), ritonavir (Norvir®, Kaletra® – used to treat HIV infection), or digoxin (Lanoxin® – used to treat heart problems).