

FOR ADULTS WITH TYPE 2 DIABETES

many people have found they can **love their numbers**—including someone you may recognize

Type 2 diabetes can affect anyone—even James Earl Jones.

Watch Mr. Jones talk about his amazing life and work, type 2 diabetes, and how

he loved his numbers with INVOKANA®.

Watch Mr. Jones' story at INVOKANA.com/Jones

Possible Side Effects of INVOKANA®

INVOKANA® may cause serious side effects, including:

- **Ketoacidosis** (increased ketones in your blood or urine). Ketoacidosis has happened in people who have type 1 or type 2 diabetes, during treatment with INVOKANA®. Ketoacidosis can be life-threatening and may need to be treated in a hospital. Ketoacidosis can happen with INVOKANA® even if your blood sugar is less than 250 mg/dL. Stop taking INVOKANA® and call your doctor right away if you get any of the following symptoms: nausea, vomiting, stomach-area pain, tiredness, or trouble breathing
- **Kidney problems.** Sudden kidney injury has happened to people taking INVOKANA®. Talk to your doctor right away if you: 1) reduce the amount of food or liquid you drink, if you are sick, or cannot eat or 2) you start to lose liquids from your body from vomiting, diarrhea, or being in the sun too long
- **A high amount of potassium in your blood (hyperkalemia)**
- **Serious Urinary Tract Infections:** may lead to hospitalization and have happened in people taking INVOKANA®. Tell your doctor if you have signs or symptoms of a urinary tract infection such as: burning feeling while urinating, need to urinate often or right away, pain in the lower part of your stomach (pelvis), or blood in the urine. Some people may also have high fever, back pain, nausea, or vomiting
- **Low blood sugar (hypoglycemia).** If you take INVOKANA® with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin, your risk of getting low blood sugar is higher. The dose of your sulfonylurea medicine or insulin may need to be lowered while you take INVOKANA®

Signs and symptoms of low blood sugar may include: headache, drowsiness, weakness, dizziness, confusion, irritability, hunger, fast heartbeat, sweating, shaking, or feeling jittery.

Serious allergic reaction. If you have any symptoms of a serious allergic reaction, stop taking INVOKANA® and call your doctor right away or go to the nearest hospital emergency room.

Broken Bones (fractures): Bone fractures have been seen in patients taking INVOKANA®. Talk to your doctor about factors that may increase your risk of bone fracture.

The most common side effects of INVOKANA® include: vaginal yeast infections and yeast infections of the penis; changes in urination, including urgent need to urinate more often, in larger amounts, or at night.

Tell your doctor if you have any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Janssen Scientific Affairs, LLC at 1-800-526-7736.

Please see Medication Guide on the next page.

Canagliflozin is licensed from Mitsubishi Tanabe Pharma Corporation.

Trademarks are those of their respective owners.

© Janssen Pharmaceuticals, Inc. 2016 September 2016 044987-160615

once-daily
Invokana®
canagliflozin tablets



imagine loving your numbers

053652-160523