

Blackboard

# SafeAssign Originality Report

NURS-6630N-27-Approaches to Treatment-2021-Su... • SafeAssign Drafts • Submitted on Sun, Jul 18, 2021, 3:28 AM

[View Originality Report - Old Design](#)

View Report Summary

On the other hand, Wellbutrin would be an appropriate option because it helps treat symptoms of seasonal affective disorder and depression. (7) However, the drug has severe side effects, including suicidal thoughts and seizures. By reducing the Zyprexa dosage from 10mg to 7.5mg, there is the expectation that the patient would not gain extra weight (Lähteenvuori et al., 2020). (4) The revised dosage of 7.5mg was also within her weight. (5) I advised the patient to actively involve in regular physical exercise, observe normal meal schedules, and stop snacking a whole day to lose the initially added excess weight. In this case, a major ethical consideration is an informed consent. Schizophrenia patients mainly have neurological deficits that make them psychologically unfit to make personal decisions concerning treatment. Again, it is challenging to understand what the patients are consenting to. Hence, I would take caution when trying to get voluntary and informed consent from the patient to ensure that no decision would harm the patient. In this case, I would strike a

|             |               |                                       |                                      |
|-------------|---------------|---------------------------------------|--------------------------------------|
| Word Count: | Submitted on: | Submission UUID:                      | Attachment UUID:                     |
| 1,821       | 07/18/21      | d411f247a-6b32-fb9e-f795-30a82b9bb154 | 776a8a42-9f5f-9d97-d955-1d628f2e324a |

## Sources

### INCLUDED SOURCES

Institutional database (7) 38 %

Global database (5) 6 %

Internet (2) 2 %