

SafeAssign Originality Report

NURS-6630N-27-Approaches to Treatment-2021-Su... • SafeAssign Drafts • Submitted on Sun, Aug 01, 2021, 10:36 AM

[View Originality Report - Old Design](#)

 [View Report Summary](#)

mental health disorders are ruled out by lack of delusional or paranoid thought processes and suicidal ideation. The condition presents predominantly in inattentive behaviour, which reflects in various forms. At eight years, the best treatment mechanism is medication. The child was put on medication with periodic evaluations to determine the progress and establish the basis for any necessary change. (4) Decision #1 The first decision was to begin Wellbutrin (bupropion) XL 150 mg orally daily for four weeks. (5) The medicine is used off-label for the treatment of ADHD but is most commonly used in adults. Wellbutrin increases dopamine neurotransmission in the frontal cortex and hence its effectiveness in treating the condition. However, the medicine had to be dropped since it cause suicidal ideation in the patient. Suicidal ideation is a common side effect on children and adolescents (Davis, 2018). This side effect is highly associated with antidepressants. The other two choices were not picked since they also caused side effects despite the slight improvement in the patient's symptoms. Intuniv extended-release produced no positive progress in the

Word Count:	Submitted on:	Submission UUID:	Attachment UUID:
1,591	08/01/21	0461a735-12ce-63ce-0219-28973d69f8a4	41716ece-27db-1674-99c3-235f28b5c5d10



Attachment 1

WK9AssgnAnimashaunO Psych...

49%

Sources

INCLUDED SOURCES

Institutional database (5)

46%

Internet (2)

3%