

SafeAssign Originality Report

NURS-6630N-27-Approaches to Treatment-2021-Su... • SafeAssign Drafts • Submitted on Sun, Aug 01, 2021, 10:36 AM

[View Originality Report - Old Design](#)

📄

View Report Summary

the disorder. Also, the doctor hoped to eliminate the side effect the drug caused in four weeks (Bachmann et al., 2017). ③ The physicians considered the ethical issues of privacy, confidentiality, safety and informed consent. The psychiatrist was careful to suggest the most effective drugs that had little or no possible side effects to ensure the safety of the patient. The girl's parents were involved in making the medication decisions after they were provided with all essential information concerning the drugs. Conclusion The eight-year-old girl presented symptoms that suggested she had ADHD. The most appropriate treatment method involved medication. ④ The treatment began with Wellbutrin (bupropion) XL 150 mg orally daily for four weeks. The drug did not cause any significant positive progress. Also, the medicine had to be dropped since it caused suicidal ideation in the patient. Instead, the drug was replaced with Strattera, which the patient would take at 10 mg daily. After four weeks of medication, the girl exhibited improvement in the symptoms of ADHD. However, the medicine also caused decreased appetite in the patient.

Word Count:	Submitted on:	Submission UUID:	Attachment UUID:
1,591	08/01/21	0461a735-12ce-63ce-0219-28973d69f8a4	41716ece-27cb-1674-99c3-235f28b5cd10

✖

Attachment 1

WK9AssgnAnimashaunO Psych...

49 %

Sources	^
INCLUDED SOURCES	🌈 📄
Institutional database (5)	46 % ⌵
Internet (2)	3 % ⌵