

The Mozart effect

J S Jenkins MD FRCP

J R Soc Med 2001;94:170-172

In 1993 Rauscher *et al.*¹ made the surprising claim that, after listening to Mozart's sonata for two pianos (K448) for 10 minutes, normal subjects showed significantly better spatial reasoning skills than after periods of listening to relaxation instructions designed to lower blood pressure or silence. The mean spatial IQ scores were 8 and 9 points higher after listening to the music than in the other two conditions. The enhancing effect did not extend beyond 10-15 minutes. These results proved controversial. Some investigators were unable to reproduce the findings²⁻⁴ but others confirmed that listening to Mozart's sonata K448 produced a small increase in spatial-temporal performance, as measured by various tests derived from the Stanford-Binet scale such as paper-cutting and folding procedures⁵⁻⁷ or pencil-and-paper maze tasks⁸. However, Rauscher has stressed that the Mozart effect is limited to spatial temporal reasoning and that there is no enhancement of general intelligence; some of the negative results, she thinks, may have been due to inappropriate test procedures⁹.

So, does the Mozart effect exist? The generality of the original positive findings has been criticized on the grounds that any Mozart effect is due to 'enjoyment arousal' occasioned by this particular music and would not take place in the absence of its appreciation. This interpretation is countered by animal experiments in which separate groups of rats were exposed, *in utero* followed by a postpartum period of 60 days, to Mozart's piano sonata K448, to minimalist music by the composer Philip Glass, to white noise or to silence and then tested for their ability to negotiate a maze. The Mozart group completed the maze test significantly more quickly and with fewer errors ($P < 0.01$) than the other three groups; thus, enjoyment and musical appreciation is unlikely to have been the basis of the improvement¹⁰.

LOCALIZATION OF MUSIC PERCEPTION AND SPATIAL IMAGING WITHIN THE BRAIN

An explanation for the results obtained after listening to music may lie in the manner in which music and spatial imaging are processed within the brain. There have been many studies on the localization of music perception. Techniques such as positron emission tomography (PET)

and functional magnetic resonance scanning, together with studies on localized brain lesions, have shown that listening to music activates a wide distribution of brain areas. The primary auditory area lies classically in the transverse and superior temporal gyri, but particular components of musical appreciation involving rhythm, pitch, metre, melody, and timbre are processed in many different areas of the brain. These range from the prefrontal cortex and superior temporal gyrus to the precuneus of the parietal lobe, with much interconnection of the different networks activated¹¹⁻¹³. Rhythm and pitch discrimination are processed mainly in the left hemisphere whereas timbre and melody are found chiefly in the right. Appreciation of metre does not appear to show hemispheric preference.

Brain areas concerned with mental imaging as tested by spatial temporal tasks (such as the building of three-dimensional cube assemblies in sequence) were also mapped by PET scanning¹⁴. The results show that the areas activated include the prefrontal, temporal and precuneus regions which overlap with those involved in music processing. It is suggested, therefore, that listening to music would prime the activation of those areas of the brain which are concerned with spatial reasoning.

LONG-TERM EFFECTS OF MUSIC ON THE BRAIN

The original experiments on adults exposed to Mozart's music were of short duration only. In related experiments¹⁵, long-term effects of music were studied in groups of pre-school children aged 3-4 years who were given keyboard music lessons for six months, during which time they studied pitch intervals, fingering techniques, sight reading, musical notation and playing from memory. At the end of training all the children were able to perform simple melodies by Beethoven and Mozart. When they did they were then subjected to spatial-temporal reasoning tests calibrated for age, and their performance was more than 30% better than that of children of similar age given either computer lessons for 6 months or no special training ($P < 0.001$). The improvement was limited to spatial-temporal reasoning; there was no effect on spatial recognition. The effect lasted unchanged for 24 hours after the end of the music lessons but the precise duration of the enhancement was not further explored. The longer duration of the effects than in previous reports was

attributed to the length of exposure to music and the greater plasticity of the young brain. In further experiments of this kind it has been claimed that the enhancement of spatial-temporal reasoning in children after piano training has resulted in significantly greater scores in higher mathematics¹⁶.

MUSIC AND THE ELECTROENCEPHALOGRAPHIC PATTERN

Attempts have been made to investigate the electrical discharge patterns of brain areas after exposure to music. In one study, listening to the Mozart sonata K448 for 10 minutes, in contrast to listening to a short story, resulted in enhanced synchrony of the firing pattern of the right frontal and left temporoparietal areas of the brain, which persisted for 12 minutes⁶. Listening to the sonata was also accompanied by increased power of the beta spectrum of the electroencephalogram in the right temporal, left temporal and right frontal regions¹⁷. In a further study, listening to music (not that of Mozart) also resulted in greater beta power, particularly in the area of the precuneus bilaterally¹⁸.

MOZART EFFECT ON EPILEPSY

A more impressive indication of a Mozart effect is to be seen in epilepsy. In 23 of 29 patients with focal discharges or bursts of generalized spike and wave complexes who listened to the Mozart piano sonata K448 there was a significant decrease in epileptiform activity as shown by the electroencephalogram (EEG)¹⁹. Some individual patients showed especially striking improvement. In one male, unconscious with status epilepticus, ictal patterns were present 62% of the time, whereas during exposure to Mozart's music this value fell to 21%. In two other patients with status epilepticus continuous bilateral spike and wave complexes were recorded 90–100% of the time before the music, suddenly falling to about 50% 5 minutes after the music began. The fact that improvement took place even in a comatose patient demonstrates again that appreciation of the music is not a necessary feature of the Mozart effect.

To determine whether this music could exert a longer effect, studies were conducted in an eight-year-old girl with a particularly intractable form of childhood epilepsy, the Lennox–Gastaut syndrome, with many drop attacks accompanied by bilateral spike and wave complexes and focal discharges from the right posterior temporal area²⁰. Mozart's sonata was played every 10 minutes for each hour of the day when she was awake. At the end of the waking period the number of clinical seizures had fallen from 9 during the initial four hours to one during the last four hours and the number of seconds during which general

discharges occurred fell from 317 to 178. The following day the number of attacks was two in seven and half hours.

SPECIFICITY OF MOZART'S MUSIC

To what extent are the changes attributable specifically to Mozart's music? Following the initial experiments of Rauscher *et al.*¹ most researchers have used Mozart's double piano sonata K448, which the Mozart authority Alfred Einstein called 'one of the most profound and most mature of all Mozart's compositions', but his piano concerto no 23 in A major K488 also proved to be effective⁸. Some investigators observed that no enhancement of spatial temporal tests was seen after the minimalist music of Philip Glass⁵, and there was no improvement in epileptiform EEG tracings after old-time pop music¹⁹. Rideout *et al.*, however, report that a contemporary composition by the Greek-American musician Yanni, which they suggest is similar to the Mozart sonata in tempo, structure, melody and harmony, was also effective⁷. In an attempt to determine the physical characteristics which were responsible for the Mozart effect, Hughes and Fino²¹ subjected a wide range of music to computer analysis. As many as 81 selections of Mozart, 67 of J C Bach, 67 of J S Bach, 39 of Chopin, and 148 from 55 other composers were analysed. The characteristic shown by much of Mozart's music and shared with the two Bachs was a high degree of long-term periodicity, especially within the 10–60 s range.

Another similarity between the music of Mozart and the two Bachs was the emphasis on the average power of particular notes, notably G3 (196 Hz), C5 (523 Hz) and B5 (987 Hz). In contrast, Philip Glass' minimalist music and old-time pop music, which had both proved without effect on spatial behavioural tasks or on epilepsy, showed little long-term periodicity. It is suggested that music with a high degree of long-term periodicity, whether of Mozart or other composers, would resonate within the brain to decrease seizure activity and to enhance spatial-temporal performance.

CONCLUSION

An enhancement of spatial-temporal reasoning performance after listening to Mozart's music for 10 minutes has been reported by several, but not all, researchers. Even in the studies with positive results the enhancement is small and lasts about 12 minutes. The effect varies between individuals and depends upon the spatial tasks chosen; general intelligence is not affected. Rather more impressively, there is a beneficial effect on some patients with epilepsy. The results are not specific to Mozart's compositions but the exact musical criteria required have not been completely defined.

The practical use of such observations is as yet uncertain, especially since many of the experiments relate only to short listening periods to Mozart's piano sonata K448. More studies are necessary, involving longer-term exposure to Mozart and to a wide selection of other composers, before the effect can be fully assessed.

REFERENCES

- 1 Rauscher FH, Shaw GL, Ky KN. Music and spatial task performance. *Nature* 1993;365:611
- 2 Chabris CF. Prelude or requiem for the 'Mozart effect'? *Nature* 1999;400:826-7
- 3 Steele KM, Dalla Bella S, Peretz I, *et al.* Prelude or requiem for the 'Mozart effect'? *Nature* 1999;400:827
- 4 Newman J, Rosenback JH, Burns IL, *et al.* An experimental test of "the Mozart effect": does listening to his music improve spatial ability? *Percept Motor Skills* 1995;81:1379-87
- 5 Rauscher FH, Shaw GL, Ky KN. Listening to Mozart enhances spatial-temporal reasoning: toward a neurophysiological basis. *Neurosci Lett* 1995;185:44-7
- 6 Rideout BE, Laubach CM. EEG correlates of enhanced spatial performance following exposure to music. *Percept Motor Skills* 1996;82:427-32
- 7 Rideout BE, Dougherty S, Wernert L. Effect of music on spatial performance: a test of generality. *Percept Motor Skills* 1998;86:512-14
- 8 Wilson TL, Brown TL. Re-examination of the effect of Mozart's music on spatial task performance. *J Psychol* 1997;131:365-70
- 9 Rauscher FH. Comment. *Nature* 1999;400:827
- 10 Rauscher FH, Robinson KD, Jens JJ. Improved maze learning through early music exposure in rats. *Neurol Res* 1998;20:427-32
- 11 Warren JD. Variations on the musical brain. *J R Soc Med* 1999;92:571-5
- 12 Platel H, Price C, Baron JC, *et al.* The structural components of music perception: a functional anatomical study. *Brain* 1997;120:229-43
- 13 Ligeois-Chauvel C, Peretz I, Babai M, *et al.* Contribution of different cortical areas in the temporal lobes to music processing. *Brain* 1998;121:1853-67
- 14 Mellet E, Tzourio N, Crivello F, *et al.* Functional anatomy of spatial imagery generated from verbal instructions. *J Neurosci* 1996;16:6504-12
- 15 Rauscher FH, Shaw GL, Levine LJ, *et al.* Music training causes long-term enhancement of preschool children's spatial-temporal reasoning. *Neurol Res* 1997;19:2-8
- 16 Graziano AB, Peterson M, Shaw GL. Enhanced learning of proportional maths through music training and spatial-temporal reasoning. *Neurol Res* 1999;21:139-52
- 17 Sarnthein J, von Stein A, Rappelsberger P, *et al.* Persistent patterns of brain activity: an EEG coherence study of the positive effect of music on spatial-temporal reasoning. *Neurol Res* 1997;19:107-16
- 18 Nakamura S, Sadato N, Oohashi T, *et al.* Analysis of music-brain interaction with simultaneous measurement of regional blood flow and electroencephalogram beta rhythm in human subjects. *Neurosci Lett* 1999;275:222-6
- 19 Hughes JR, Daaboul Y, Fino JJ, *et al.* The Mozart effect on epileptiform activity. *Clin Electroencephalogr* 1998;29:109-19
- 20 Hughes JR, Fino JJ, Melyn MA. Is there a chronic change of the "Mozart effect" on epileptiform activity? A case study. *Clin Electroencephalogr* 1999;30:44-5
- 21 Hughes JR, Fino JJ. The Mozart effect: distinctive aspects of the music—a clue to brain coding? *Clin Electroencephalogr* 2000;31:94-103

Copyright of Journal of the Royal Society of Medicine is the property of Royal Society of Medicine Press Limited and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.

SCIENTIFIC CORRESPONDENCE

Music and spatial task performance

SIR—There are correlational¹, historical² and anecdotal³ relationships between music cognition and other 'higher brain functions', but no causal relationship has been demonstrated between music cognition and cognitions pertaining to abstract operations such as mathematical or spatial reasoning. We performed an

spatial IQ scores of 119, 111 and 110, respectively. Thus, the IQs of subjects participating in the music condition were 8–9 points above their IQ scores in the other two conditions. A one-factor (listening condition) repeated measures analysis of variance (ANOVA) performed on SAS revealed that subjects performed better on the abstract/spatial reasoning tests after listening to Mozart than after listening to either the relaxation tape or to nothing ($F_{2,35} = 7.08$; $P = 0.002$). The music condition differed significantly from both the relaxation and the silence conditions (Scheffe's $t = 3.41$, $P = 0.002$; $t = 3.67$, $P = 0.0008$, two-tailed, respectively).

The relaxation and silence conditions did not differ ($t = 0.795$; $P = 0.432$, two-tailed). Pulse rates were taken before and after each listening condition. A two-factor (listening condition and time of pulse measure) repeated measures ANOVA revealed no interaction or main effects for pulse, thereby excluding arousal as an obvious cause. We found no order effects for either condition presentation or task, nor any experimenter effect.

The enhancing effect of the music condition is temporal, and does not extend beyond the 10–15-minute period during which subjects were engaged in each spatial task. Inclusion of a delay period (as a variable) between the music listening condition and the testing period would allow us quantitatively to determine the presence of a decay constant. It would also be interesting to vary the listening time to optimize the enhancing effect, and to examine whether other measures of general intelligence (verbal reasoning, quantitative reasoning and short-term memory) would be similarly facilitated. Because we used only one musical sample of one composer, various other compositions and musical styles

should also be examined. We predict that music lacking complexity or which is repetitive may interfere with, rather than enhance, abstract reasoning. Also, as musicians may process music in a different way from non-musicians, it would be interesting to compare these two groups.

Frances H. Rauscher


Gordon L. Shaw*

Katherine N. Ky

Center for the Neurobiology of Learning
 and Memory,
 University of California,
 Irvine, California 92717, USA

* Also at Department of Physics.

1. Hassler, M., Birbaumer, N. & Feil, A. *Psychol. Music* **13**, 99–113 (1985).
2. Allman, G. J. *Greek Geometry from Thales to Euclid* p. 23 (Arno, New York, 1976).
3. Cranberg, L. D. & Albert, M. L. in *The Exceptional Brain* (eds Obler, L. K. & Fein, D.) 156 (Guilford, New York, 1988).
4. Thorndike, R. L., Hagen, E. P. & Sattler, J. M. *The Stanford-Binet Scale of Intelligence* (Riverside, Chicago, 1986).



Standard age scores for each of the three listening conditions.

Testing procedure. In the music condition, the subject listened to 10 min of the Mozart piece. The relaxation condition required the subject to listen to 10 min of relaxation instructions designed to lower blood pressure. The silence condition required the subject to sit in silence for 10 min. One of three abstract reasoning tests taken from the Stanford-Binet intelligence scale⁴ was given after each of the listening conditions. The abstract/spatial reasoning tasks consisted of a pattern analysis test, a multiple-choice matrices test and a multiple-choice paper-folding and cutting test. For our sample, these three tasks correlated at the 0.01 level of significance. We were thus able to treat them as equal measures of abstract reasoning ability.

Scoring. Raw scores were calculated by subtracting the number of items failed from the highest item number administered. These were then converted to SAS using the Stanford-Binet's SAS conversion table of normalized standard scores with a mean set at 50 and a standard deviation of 8. IQ equivalents were calculated by first multiplying each SAS by 3 (the number of subtests required by the Stanford-Binet for calculating IQs). We then used their area score conversion table, designed to have a mean of 100 and a standard deviation of 16, to obtain SAS IQ equivalents.

experiment in which students were each given three sets of standard IQ spatial reasoning tasks; each task was preceded by 10 minutes of (1) listening to Mozart's sonata for two pianos in D major, K488; (2) listening to a relaxation tape; or (3) silence. Performance was improved for those tasks immediately following the first condition compared to the second two.

Thirty-six college students participated in all three listening conditions. Immediately following each listening condition, the student's spatial reasoning skills were tested using the Stanford-Binet intelligence scale⁴. The mean standard age scores (SAS) for the three listening conditions are shown in the figure. The music condition yielded a mean SAS of 57.56; the mean SAS for the relaxation condition was 54.61 and the mean score for the silent condition was 54.00. To assess the impact of these scores, we 'translated' them to

delay period (as a variable) between the music listening condition and the testing period would allow us quantitatively to determine the presence of a decay constant. It would also be interesting to vary the listening time to optimize the enhancing effect, and to examine whether other measures of general intelligence (verbal reasoning, quantitative reasoning and short-term memory) would be similarly facilitated. Because we used only one musical sample of one composer, various other compositions and musical styles

Scientific Correspondence

Scientific Correspondence is intended to provide a forum in which readers may raise points of a scientific character. They need not arise out of anything published in *Nature*. In any case, priority will be given to letters of fewer than 500 words and five references.

MyoD and c-fos expression

SIR—Trousche *et al.*¹ have reported that the down regulation of *c-fos* expression during muscle cell differentiation may result from the binding of the helix-loop-helix (HLH) proteins to a CANNTG motif, or E-box, that occurs within the *c-fos* serum response element (SRE), thereby excluding the binding of the serum response factor (SRF) to this element. We investigated the interaction of HLH proteins with the *c-fos* SRE when molecular clones for E12 were first isolated by screening a phage expression library with the SRE probe². We estimated that the dissociation constant for the myogenin/E12-SRE complex was 10^{-8} – 10^{-9} M⁻¹ by comparing the relative effectiveness of E-box elements from different genes to compete for binding in the electrophoretic mobility-shift assays. The relatively low affinity of myogenin/E12 for the SRE could result from differences in nucleotide identities at the internal dinucleotide and flanking sequences between the *c-fos* E-box and the consensus HLH binding site^{3,4}. By contrast, the dissociation constants for the SRE-SRF or the SRE-SRF/p62^{TCF} complexes have been reported to be 5×10^{-10} and $\leq 10^{-11}$ M⁻¹, respectively⁵. Collectively, these data indicate that it is unlikely that HLH proteins alone can significantly compete with SRF for binding to DNA *in vivo*.

It has been reported in other studies that the *c-fos* SRE is either equally active in muscle and non-muscle cells⁶, or that it activates muscle-specific expression⁷ when situated upstream from a minimal promoter. Further, a comparison of the